

Operations Research, Spring 2017
Lecture 9: Single-variate Nonlinear Programming

Instructor: Ling-Chieh Kung
Department of Information Management
National Taiwan University

1. (a) What is the maximum area inside a rectangle whose lengths of the four edges sum to 1? Formulate an NLP whose optimal solution answers this question.
- (b) What is the maximum total length of a rectangle whose area is 1? Formulate an NLP whose optimal solution answers this question.

2. Graphically or intuitively determine which of the following sets are convex:

(a) An interval $[a, b] \subseteq \mathbb{R}$ for some $a, b \in \mathbb{R}$.

(b) $[0, 10] \cup [20, 30]$.

(c) $\{x \in \mathbb{R}^2 \mid x_1 + x_2 \leq 2, x_1^2 + x_2^2 \leq 9\}$.

(d) $\{x \in \mathbb{R}^3 \mid x_1 + 2x_2 + 4x_3 \leq 8, x_2 \geq 0, x_3 \geq 0\}$.

(e) $\{x \in \mathbb{Z}^2 \mid x_1 \geq 0, x_2 \geq 0, x_1 + 2x_2 \leq 6\}$.

3. Graphically or intuitively determine which of the following functions are convex over the given domain S :

(a) $f(x) = x^3$, $S = \mathbb{R}$.

(b) $f(x) = x^3$, $S = [0, \infty)$.

(c) $f(x) = \frac{1}{x}$, $S = (0, \infty)$.

(d) $f(x) = x^a$ for some $a \in (0, 1)$, $S = [0, \infty)$.

(e) $f(x) = x^a$ for some $a \in (1, 2)$, $S = [0, \infty)$.

(f) $f(x) = 2^x$, $S = \mathbb{R}$.

4. For each of the following functions over given domains, graphically find all local and global minima.

(a) $f(x) = x^3 + 2x^2 - 2$ over $[-2, 2]$.

(b) $f(x) = -x^2$ over $(-1, 0] \cup [1, 2]$.

(c) $f(x) = e^x$ over \mathbb{R} .

5. For each of the following single-variate twice differentiable functions, analytically determine whether it is convex, concave, or neither in the given domain:

(a) $f(x) = x(x - 1)(x + 2)$ over $(-\infty, -1]$.

(b) $f(x) = e^x + x^3$ over $[0, \infty)$.

(c) $f(x) = \frac{1}{x} + x$ over $(0, \infty)$.

6. The problem for finding the maximum area inside a rectangle whose lengths of the four edges sum to 1 can be formulated as follows. Let x and y be the height and width of the rectangle, the formulation is

$$\begin{aligned} \max_{x \geq 0, y \geq 0} \quad & xy \\ \text{s.t.} \quad & x + y = \frac{1}{2}. \end{aligned}$$

- (a) Is the feasible region convex?
(b) Explain why the NLP is equivalent to the one below:

$$\begin{aligned} \max \quad & x \left(\frac{1}{2} - x \right) \\ \text{s.t.} \quad & 0 \leq x \leq \frac{1}{2}. \end{aligned}$$

- (c) Solve the above single-variate NLP and find an optimal solution.

7. A retailer is importing a product from an overseas supplier. If there are q units on the market, the unit price of the product will be $a - bq$ dollars (this is called the *market clearing price*). The unit procurement cost of the product is $\$c$. The retailer wants to determine a procurement quantity that maximizes its profit.
- (a) Formulate an NLP that maximizes the retailer's profit.
 - (b) Solve the NLP.
 - (c) Determine how a , b , and c affect the optimal procurement quantity and the associated profit.
 - (d) Provide economic interpretations for your answers above.

8. Each month, a gas station sells 4,000 gallons of gasoline. Each time the parent company refills the station's tanks, it charges the station a fixed cost \$50 plus a variable cost \$0.7 per gallon. The annual cost of holding a gallon of gasoline is \$0.3. Suppose the demand rate is constant and no shortage is allowed.
- (a) How large should the station's one order be?
 - (b) How many orders per year will be placed in average?
 - (c) How long will it be between orders (how long is the cycle time)?
 - (d) Suppose that there is an *ordering lead* $L > 0$, which is the amount of time it takes to get the ordered gasoline after an order is placed. Suppose that $L \leq T^*$, where T^* is the optimal cycle time. Explain why the optimal *reorder point* is to order when the on-hand inventory level is LD .
 - (e) If the lead time is a half month, what is the reorder point?
 - (f) If the lead time is 2.2 months, what is the reorder point?