

Homework Assignment #1

Note

This assignment is due 2:10PM Tuesday, March 5, 2013. Please write or type your answers on A4 (or similar size) paper. Drop your homework by the due time in Yih-Kuen Tsay's mail box on the first floor of Management College Building II. Late submission will be penalized by 20% for each working day overdue. You may discuss the problems with others, but copying answers is strictly forbidden.

Problems

There are five problems in this assignment, each accounting for 20 points. You must use *induction* for all proofs. (Note: problems marked with "(X.XX)" are taken from [Manber 1989] with probable adaptation.)

- (2.10) Find an expression for the sum of the i -th row of the following triangle, which is called the **Pascal triangle**, and prove the correctness of your claim. The sides of the triangle are 1s, and each other entry is the sum of the two entries immediately above it.

$$\begin{array}{cccccc}
 & & & & & 1 \\
 & & & & 1 & 1 \\
 & & 1 & 2 & 1 & \\
 & 1 & 3 & 3 & 1 & \\
 1 & 4 & 6 & 4 & 1 &
 \end{array}$$

- The Harmonic series $H(k)$ is defined by $H(k) = 1 + \frac{1}{2} + \frac{1}{3} + \dots + \frac{1}{k-1} + \frac{1}{k}$. Prove that $H(2^n) \geq 1 + \frac{n}{2}$, for all $n \geq 0$ (which implies that $H(k)$ diverges).
- (2.14) Consider the following series: 1, 2, 3, 4, 5, 10, 20, 40, ..., which starts as an arithmetic series, but after the first 5 terms becomes a geometric series. Prove that any positive integer can be written as a sum of distinct numbers from this series.
- (2.19) Prove that the regions formed by n circles in the plane can be colored with two colors such that any neighboring regions (which share an arc, not just a point) are colored differently.
- (2.37) Consider the recurrence relation for Fibonacci numbers $F(n) = F(n-1) + F(n-2)$. Without solving this recurrence, compare $F(n)$ to $G(n)$ defined by the recurrence $G(n) = G(n-1) + G(n-2) + 1$. It seems obvious that $G(n) > F(n)$ (because of the extra 1). Yet the following is a seemingly valid proof (by induction) that $G(n) = F(n) - 1$. We assume, by induction, that $G(k) = F(k) - 1$ for all k such that $1 \leq k \leq n$, and we consider $G(n+1)$:

$$G(n+1) = G(n) + G(n-1) + 1 = F(n) - 1 + F(n-1) - 1 + 1 = F(n+1) - 1$$

What is wrong with this proof?