

Suggested Solutions for Homework Assignment #1

We assume the binding powers of the logical connectives and the entailment symbol decrease in this order: \neg , $\{\wedge, \vee\}$, \rightarrow , \leftrightarrow , \vdash .

- (30 points) Prove that every propositional formula has an equivalent formula in the conjunctive normal form and also an equivalent formula in the disjunctive normal form. (Hint: by induction on the structure of a formula, dealing with both cases simultaneously)

Solution. Let us first review some preliminaries. A (non-empty) clause is a disjunction of one or more literals such as $p \vee \neg q \vee r$, while a (non-empty) term/product is a conjunction of one or more literals such as $\neg p \wedge q \wedge \neg r$. (Note: the name “term” as defined here is not commonly used in propositional logic. However, it is adequate in light of the notion of a term in algebraic expressions. An alternative name is “product”.) So, a formula is in conjunctive normal form (CNF) if it is a conjunction of one or more clauses. A formula is in disjunctive normal form (DNF) if it is a disjunction of one or more terms. A clause by itself is in CNF (a one-clause CNF) and, when seen as a disjunction of one-literal terms, is also in DNF. Similarly, for a term. A single literal is a special case of a clause and also of a term.

The complement of a clause (term), after the negation is pushed to the literal level, becomes a term (clause), e.g., $\neg(p \vee \neg q \vee r) \Leftrightarrow \neg p \wedge q \wedge \neg r$. Taking this one level up, the complement of a formula in CNF (DNF), after the negation is pushed to the literal level, becomes a formula in DNF (CNF), e.g., $\neg((p \vee \neg q) \wedge (q \vee r)) \Leftrightarrow (\neg p \wedge q) \vee (\neg q \wedge \neg r)$.

Now we prove the problem statement by induction on the structure of a given formula φ .

Base case (φ is just a propositional symbol): a propositional symbol can be seen as a single-literal clause or term and so is already in CNF and in DNF.

Inductive step: there are three cases.

- $\varphi = \neg\psi$: let ψ^C be a formula equivalent to ψ in CNF and ψ^D an equivalent formula in DNF (guaranteed to exist by the induction hypothesis). Pushing the negation at the front of $\neg\psi^C$ ($\neg\psi^D$) to the literal level, we get a formula equivalent to φ in DNF (CNF).
- $\varphi = \varphi_1 \wedge \varphi_2$: let φ_1^C (φ_2^C) be a formula equivalent to φ_1 (φ_2) in CNF and φ_1^D (φ_2^D) an equivalent formula in DNF. The formula $\varphi_1^C \wedge \varphi_2^C$ is equivalent to φ and readily in CNF.

To obtain a formula equivalent to φ in DNF, suppose $\varphi_1^D = t_1 \vee t_2 \vee \dots \vee t_l$ and $\varphi_2^D = u_1 \vee u_2 \vee \dots \vee u_m$, where t_i 's and u_j 's are terms. Then, by repeatedly distributing the top-level \wedge in $\varphi_1^D \wedge \varphi_2^D$ to the term level, we obtain a formula $\bigvee_{1 \leq i \leq l, 1 \leq j \leq m} (t_i \wedge u_j)$ in DNF that is equivalent to φ .

- $\varphi = \varphi_1 \vee \varphi_2$: analogous to the case of $\varphi = \varphi_1 \wedge \varphi_2$.

□

- (40 points) Prove, using *Natural Deduction* (in the sequent form), the validity of the following sequents:

(a) $\vdash (p \rightarrow (q \rightarrow r)) \rightarrow (p \wedge q \rightarrow r)$

Solution.

$$\frac{\frac{\frac{\frac{\alpha}{p \rightarrow (q \rightarrow r), p \wedge q \vdash q} (\rightarrow E)}{p \rightarrow (q \rightarrow r), p \wedge q \vdash r} (\rightarrow I)}{p \rightarrow (q \rightarrow r) \vdash p \wedge q \rightarrow r} (\rightarrow I)}{\vdash (p \rightarrow (q \rightarrow r)) \rightarrow (p \wedge q \rightarrow r)} (\rightarrow I)$$

α :

$$\frac{\frac{\frac{\alpha}{p \rightarrow (q \rightarrow r), p \wedge q \vdash p \rightarrow (q \rightarrow r)} (Hyp)}{p \rightarrow (q \rightarrow r), p \wedge q \vdash q \rightarrow r} (\rightarrow E)}{\frac{\frac{\frac{\frac{\alpha}{p \rightarrow (q \rightarrow r), p \wedge q \vdash p \wedge q} (\wedge E_2)}{p \rightarrow (q \rightarrow r), p \wedge q \vdash p \wedge q} (Hyp)}{p \rightarrow (q \rightarrow r), p \wedge q \vdash p} (\wedge E_1)}{p \rightarrow (q \rightarrow r), p \wedge q \vdash q \rightarrow r} (\rightarrow E)}$$

□

(b) $\vdash (p \wedge q \rightarrow r) \rightarrow (p \rightarrow (q \rightarrow r))$

Solution.

$$\frac{\frac{\frac{\frac{\frac{\alpha}{p \wedge q \rightarrow r, p, q \vdash p \wedge q \rightarrow r} (Hyp)}{p \wedge q \rightarrow r, p, q \vdash p} (\rightarrow I)}{p \wedge q \rightarrow r, p, q \vdash q \rightarrow r} (\rightarrow I)}{\vdash (p \wedge q \rightarrow r) \rightarrow (p \rightarrow (q \rightarrow r))} (\rightarrow I)}{\frac{\frac{\frac{\frac{\frac{\alpha}{p \wedge q \rightarrow r, p, q \vdash p} (Hyp)}{p \wedge q \rightarrow r, p, q \vdash p \wedge q} (\wedge I)}{p \wedge q \rightarrow r, p, q \vdash p \wedge q} (\rightarrow E)}{p \wedge q \rightarrow r, p, q \vdash p} (\rightarrow I)}{p \wedge q \rightarrow r, p, q \vdash q \rightarrow r} (\rightarrow I)}{p \wedge q \rightarrow r, p, q \vdash p \wedge q \rightarrow r} (\wedge I)}$$

□

3. (30 points) Prove, using *Natural Deduction* (in the sequent form), the validity of the following sequents:

(a) $\vdash p \vee \neg p$

Solution.

$$\frac{\frac{\frac{\frac{\alpha}{\neg(p \vee \neg p) \vdash \neg p} (\neg I)}{\neg(p \vee \neg p) \vdash p \vee \neg p} (\vee I_2)}{\neg(p \vee \neg p) \vdash (p \vee \neg p) \wedge \neg(p \vee \neg p)} (\wedge I)}{\vdash \neg \neg(p \vee \neg p)} (\neg \neg E)}{\vdash p \vee \neg p} (\neg E)$$

α :

$$\frac{\frac{\frac{\frac{\alpha}{\neg(p \vee \neg p), p \vdash p} (Hyp)}{\neg(p \vee \neg p), p \vdash p \vee \neg p} (\vee I_1)}{\neg(p \vee \neg p), p \vdash (p \vee \neg p) \wedge \neg(p \vee \neg p)} (\wedge I)}{\neg(p \vee \neg p), p \vdash \neg(p \vee \neg p)} (Hyp)}$$

□

(b) $\vdash ((p \rightarrow q) \rightarrow p) \rightarrow p$ *Solution.*

$$\begin{array}{c}
\frac{}{(p \rightarrow q) \rightarrow p, \neg p \vdash (p \rightarrow q) \rightarrow p} \text{ (Hyp)} \quad \alpha \quad \frac{}{(p \rightarrow q) \rightarrow p, \neg p \vdash \neg p} \text{ (Hyp)} \\
\frac{}{(p \rightarrow q) \rightarrow p, \neg p \vdash p} \text{ (}\rightarrow E\text{)} \quad \frac{}{(p \rightarrow q) \rightarrow p, \neg p \vdash \neg p} \text{ (}\wedge I\text{)} \\
\frac{}{(p \rightarrow q) \rightarrow p, \neg p \vdash p \wedge \neg p} \text{ (}\neg I\text{)} \\
\frac{}{(p \rightarrow q) \rightarrow p \vdash \neg \neg p} \text{ (}\neg\neg E\text{)} \\
\frac{}{(p \rightarrow q) \rightarrow p \vdash p} \text{ (}\neg\neg I\text{)} \\
\frac{}{\vdash ((p \rightarrow q) \rightarrow p) \rightarrow p} \text{ (}\rightarrow I\text{)}
\end{array}$$

 $\alpha :$

$$\begin{array}{c}
\frac{}{(p \rightarrow q) \rightarrow p, \neg p, p, \neg q \vdash p} \text{ (Hyp)} \quad \frac{}{(p \rightarrow q) \rightarrow p, \neg p, p, \neg q \vdash \neg p} \text{ (Hyp)} \\
\frac{}{(p \rightarrow q) \rightarrow p, \neg p, p \vdash q} \text{ (}\rightarrow I\text{)} \quad \frac{}{(p \rightarrow q) \rightarrow p, \neg p, p, \neg q \vdash \neg p} \text{ (}\neg E\text{)} \\
\frac{}{(p \rightarrow q) \rightarrow p, \neg p \vdash p \rightarrow q} \text{ (}\rightarrow I\text{)}
\end{array}$$

□